

Eccentricity Fluctuation in Initial Conditions of Hydrodynamics

Tetsufumi Hirano^a and Yasushi Nara^b

^a*Department of Physics, The University of Tokyo, Tokyo 113-0033, Japan*

^b*Akita International University, Yuwa, Akita-city, 010-1292, Japan*

Abstract

We study effects of eccentricity fluctuations on the elliptic flow coefficient v_2 at mid-rapidity in both Au+Au and Cu+Cu collisions at $\sqrt{s_{NN}} = 200$ GeV by using a hybrid model that combines ideal hydrodynamics for space-time evolution of the quark gluon plasma phase and a hadronic transport model for the hadronic matter. We find that the effect of eccentricity fluctuation is modest in semicentral Au+Au collisions but significantly enhances v_2 in Cu+Cu collisions.

1. Introduction

One of the major discoveries at Relativistic Heavy Ion Collider (RHIC) in Brookhaven National Laboratory is that the elliptic flow [1] is found to be as large as an ideal hydrodynamic prediction for the first time in relativistic heavy ion collisions [2]. Systematic studies revealed, however, that the reasonable agreement between results from ideal hydrodynamics and elliptic flow data has been achieved only by a particular combination of dynamical modeling [3]. If one looks at the comparison carefully, one finds the agreement between hydrodynamic results and the data is not perfect. This would be due to an absence of initial fluctuation effects [4]. Therefore, further investigation is indispensable toward better understanding of the elliptic flow data and the quark gluon plasma (QGP).

2. Initial Conditions

For initial conditions in this study, we employ the Monte-Carlo version of both the Glauber model [5] and the factorized Kharzeev-Levin-Nardi (fKLN) model [6] to generate the initial distribution of entropy density on an event-by-event basis. We first calculate a transverse entropy density profile $s_0(\mathbf{x}_\perp) = s(\tau = \tau_0, x, y, \eta_s = 0)$ in each sample at an impact parameter b for a given centrality, where $\tau_0 = 0.6$ fm/c is the initial time for the hydrodynamical simulations. Then the variances of the profile are obtained from

$$\sigma_x^2 = \langle x^2 \rangle - \langle x \rangle^2, \quad (1)$$

$$\sigma_y^2 = \langle y^2 \rangle - \langle y \rangle^2, \quad (2)$$

$$\sigma_{xy} = \langle xy \rangle - \langle x \rangle \langle y \rangle. \quad (3)$$

Here $\langle \dots \rangle$ describes the average over transverse plane by weighting the entropy density in a single sample:

$$\langle \dots \rangle = \frac{\int d^2x_\perp \dots s_0(\mathbf{x}_\perp)}{\int d^2x_\perp s_0(\mathbf{x}_\perp)}. \quad (4)$$

The eccentricity with respect to the reaction plane, the participant eccentricity, and the corresponding transverse areas can be defined [7], respectively, as

$$\varepsilon_{\text{RP}} = \frac{\sigma_y^2 - \sigma_x^2}{\sigma_y^2 + \sigma_x^2}, \quad \varepsilon_{\text{part}} = \frac{\sqrt{(\sigma_y^2 - \sigma_x^2)^2 + 4\sigma_{xy}^2}}{\sigma_y^2 + \sigma_x^2}, \quad (5)$$

$$S_{\text{RP}} = \pi \sqrt{\sigma_x^2 \sigma_y^2}, \quad S_{\text{part}} = \pi \sqrt{\sigma_x^2 \sigma_y^2 - \sigma_{xy}^2}. \quad (6)$$

The impact parameter vector and the true reaction plane are not known experimentally. So one can set an apparent frame of created matter shifted by $(x, y) = (\langle x \rangle, \langle y \rangle)$ and tilted by Ψ from the true frame in the transverse plane [7]:

$$\tan 2\Psi = \frac{\sigma_y^2 - \sigma_x^2}{2\sigma_{xy}}. \quad (7)$$

The anisotropy of particle distribution could be correlated with the misidentified frame. To account for this, we first shift the center-of-mass of the system to the origin in the calculation frame and then rotate the profile in the azimuthal direction by Ψ to match the apparent reaction plane to the true reaction plane. We generate the next sample of an entropy density profile again as above and average the profiles over many samples. We repeat the above procedure for many samples until the initial distribution is smooth enough. The initial conditions obtained in this way contain the effects of eccentricity fluctuation even though the profile is smooth. In particular, even in case of vanishing impact parameter, the eccentricity is finite due to its event-by-event fluctuation. It is the particle distribution calculated from the initial conditions mentioned above that can be directly compared with the experimental data. Note that the procedure of averaging over many samples without shift or rotation corresponds to conventional initialization without the effect of eccentricity fluctuation.

By using the initial conditions above, we simulate space-time dynamics of matter produced in relativistic heavy ion collisions with a hybrid approach in which the macroscopic description of the QGP is followed by the microscopic description of the hadron gas. For details, see Ref. [8].




Figure 1: Eccentricity as a function of N_{part} in (a) Au+Au and (b) Cu+Cu collisions. Solid, dotted, dashed and dash-dotted lines correspond to $\varepsilon_{\text{part}}$ from the MC-KLN model, $\varepsilon_{\text{part}}$ from the MC-Glauber model, ε_{RP} from the MC-KLN model and ε_{RP} from the MC-Glauber model, respectively.

3. Results

Figure 1 shows the eccentricities as functions of the number of participants N_{part} with or without eccentricity fluctuations. The results from the MC-Glauber and MC-KLN models are compared with each other in (a) Au+Au and (b) Cu+Cu collisions. In semicentral Au+Au collisions at 10-50% centrality, the effect of initial fluctuations enhances the eccentricity parameter by 8-11% (5-8%) in the MC-Glauber (MC-KLN) model. The enhancement factor $\varepsilon_{\text{part}}/\varepsilon_{\text{RP}}$ is the largest at the very central bin (0-5% centrality): $\varepsilon_{\text{part}}/\varepsilon_{\text{RP}} = 1.83$ in the MC-Glauber model and 1.53 in the MC-KLN model. A qualitatively similar behavior is observed in Cu+Cu collisions, but quantitatively the effect of fluctuation is significant due to the smallness of the system.




Figure 2: Centrality dependences of v_2 for charged particles at mid-rapidity in (a) Au+Au and (b) Cu+Cu collisions at $\sqrt{s_{NN}} = 200$ GeV with the MC-Glauber model initial conditions are compared with PHOBOS data (filled plots) [9]. Open circles (squares) are results with (without) eccentricity fluctuation.




Figure 3: Centrality dependences of v_2 for charged particles at mid-rapidity in (a) Au+Au and (b) Cu+Cu collisions at $\sqrt{s_{NN}} = 200$ GeV with the MC-KLN model initial conditions are compared with PHOBOS data (filled plots) [9]. Open circles (squares) are results with (without) eccentricity fluctuation.

Figures 2 and 3 show the centrality dependence of v_2 for charged hadrons at mid-rapidity ($|\eta| < 1$) in Au+Au and Cu+Cu collisions at $\sqrt{s_{NN}} = 200$ GeV. From Figs. 2 (a) and (b), one sees that the effect of eccentricity fluctuations is not significant in Au+Au collisions, whereas v_2 is largely enhanced in Cu+Cu collisions due to fluctuation effects. These tendencies are also

expected from the results of initial eccentricity as shown in Fig. 1. It is interesting to point out that elliptic flow coefficients v_2 in the Glauber model initial conditions still slightly undershoot the experimental data, particularly in Cu+Cu collisions, even with fluctuation effects. Thus there is almost no room for the viscosity in the QGP stage to play a role in the Glauber initial conditions within the hybrid approach with the ideal gas equation of state in the QGP phase. Figure 3 is the same as Fig. 2 but the initial conditions are taken from the MC-KLN model. Again, a similar qualitative behavior is seen: The effect of eccentricity fluctuations is small in Au+Au collisions but is large in Cu+Cu collisions. Due to larger initial eccentricity in the MC-KLN model than the MC-Glauber model, the results are somewhat larger than the experimental data in peripheral Au+Au collisions, whereas we reasonably reproduce the v_2 data with initial fluctuation effects in Cu+Cu collisions. Viscous effects in the QGP phase could reduce the v_2 and enable us to reproduce the data in Au+Au collisions in this case. However, the results are already comparable with the data in Cu+Cu collisions even though the number of participants is almost the same as that in peripheral Au+Au collisions. So it would be non-trivial to establish whether the same viscous effects also give the right amount of v_2 in Cu+Cu collisions.

4. Conclusions

We calculated the elliptic flow coefficient as a function of the number of participants in the QGP hydro plus the hadronic cascade model and found that the effect of eccentricity fluctuations is visible in very central and peripheral Au+Au collisions and is quite large in Cu+Cu collisions. This strongly suggests that the effect of eccentricity fluctuations is an important factor which has to be included in the dynamical model for understanding of the elliptic flow data and for precise extraction of transport properties of the produced matter from the data.

So far, one has been focusing on comparison of hydrodynamic results with v_2 data only in Au+Au collisions. The experimental data in Cu+Cu collisions also have a strong power to constrain the dynamical models. Therefore, simultaneous analysis of v_2 data in both Au+Au and Cu+Cu collisions will be called for in future hydrodynamic studies.

Acknowledgments

The work of T.H. was partly supported by Grant-in-Aid for Scientific Research No. 19740130 and by Sumitomo Foundation No. 080734. The work of Y.N. was supported by Grand-in-Aid for Scientific Research No. 20540276.

References

- [1] J. Y. Ollitrault, Phys. Rev. D 46 (1992) 229.
- [2] T. Hirano, N. van der Kolk, A. Bilandzic, arXiv:0808.2684 [nucl-th];
- [3] T. Hirano, M. Gyulassy, Nucl. Phys. A 769 (2006) 71.
- [4] M. Miller, R. Snellings, nucl-ex/0312008.
- [5] M. L. Miller, K. Reygers, S. J. Sanders, P. Steinberg, Ann. Rev. Nucl. Part. Sci. 57 (2007) 205.
- [6] H. J. Drescher, Y. Nara, Phys. Rev. C 75 (2007) 034905;
H. J. Drescher, Y. Nara, Phys. Rev. C 76 (2007) 041903.
- [7] B. Alver *et al.* [PHOBOS Collaboration], nucl-ex/0702036;
B. Alver *et al.*, Phys. Rev. C 77 (2008) 014906.
- [8] T. Hirano, U. Heinz, D. Kharzeev, R. Lacey, Y. Nara, Phys. Lett. B 636 (2006) 299;
T. Hirano, U. Heinz, D. Kharzeev, R. Lacey, Y. Nara, Phys. Rev. C 77 (2008) 044909.
- [9] B. B. Back *et al.* [PHOBOS Collaboration], Phys. Rev. C 72 (2005) 051901;
B. Alver *et al.* [PHOBOS Collaboration], Phys. Rev. Lett. 98 (2007) 242302.